

Estimation of Incidental Interactions with Sea Turtles, Seabirds, and Marine Mammals in the 2006 Hawaii Longline Deep Set Fishery¹

Marti L. McCracken
Pacific Islands Fisheries Science Center
National Marine Fisheries Service

This report provides estimates of the number of incidental interactions with protected species by the Hawaii longline deep set fishery in the year 2006 (Table 1). Within this report, an incidental interaction means an event during a longline fishing operation in which a protected animal is hooked or entangled by the fishing gear. An incidental interaction estimate refers to the estimated total number of incidental interactions for all longline deep set fishing trips landing in the specified time period. A longline deep set fishing trip is defined as any commercial fishing trip by a vessel with a Hawaii longline permit that departs or returns at a Hawaii port, excluding those trips using a certificate for swordfishing.

The interaction estimates are based on a random sample of longline trips on which scientific observers are deployed. In 2006, observed trips were selected using two sampling schemes to accommodate fluctuating coverage levels and utilize observers efficiently. Coverage levels vary throughout the year because of fluctuation in the fleet's activity level, demands of 100% coverage in the Hawaii longline shallow set fishery, and an influx of observers after completion of NMFS observer training. Because observers are not paid while waiting to be deployed, they must be assigned with minimal delay when available. The alternative of paying them while they are waiting to be deployed would increase the cost of the observer program. The two sampling schemes attempt to reach a balance between obtaining a probability sample and being cost effective. A probability sample implies that all trips have a probability of being sampled and the sampling probabilities are known. These sampling probabilities form the basis of design-based estimators. An unbiased design-based estimator has the merit that it is unbiased regardless of the characteristics of the population being surveyed.

The primary scheme was a systematic sample. Before departing on a fishing trip, longline vessels were required to call the NOAA Fisheries Pacific Islands Regional Office (PIRO) observer program contractor at least 72 hours prior to their intended departure date. To enable sample selection, the PIRO contractor numbered calls sequentially in the order in which they were received. Herein, this assigned number is referred to as the call number. Prior to the beginning of a quarter, a systematic sample of call numbers was drawn by PIFSC and supplied to the contractor. The trips associated with these selected call numbers were designated to be sampled. Although every reasonable effort was made to sample selected trips, there were some selected trips that departed without an observer. In this situation, the PIRO contractor recorded that the trip was not sampled along with a short explanation of why it was not sampled. If a trip was selected but did not leave within a reasonable amount of time, usually the observer was reassigned to a different trip. When the selected vessel was ready to depart an observer was assigned to it.

¹ PIFSC Internal Report IR-07-006
Issued 20 April 2007

The systematic sample requires having an observer available to be deployed whenever a selected trip is ready to depart. To achieve this requirement under full targeted coverage throughout the year requires increasing the current number of observers on contract and paying them when they are not deployed. These requirements cannot be met under the current level of funding; therefore, the quarterly sample selected under the systematic design was slightly smaller than the targeted coverage, typically 5% less. The additional trips needed to reach the full targeted level were then selected using a secondary sampling scheme. This secondary scheme was used when all trips selected by the systematic sample were already covered and an observer was ready to be deployed. In this instance, a trip was randomly selected with equal probability from the calls received that day that had not already been selected. If more than one observer needed to be assigned, the appropriate number of trips was sampled with equal probability from this pool of call-ins. The coverage obtained by this secondary sampling scheme was flexible and dependent on the need to deploy observers. The additional samples drawn under the secondary sampling scheme depart from traditional probability samples because the days when additional samples were drawn were not randomly selected but determined by the need to deploy observers.

Because the systematic sample was selected quarterly, point estimates of incidental interactions were computed on a quarterly basis and then summed to estimate the year's total interactions for non-experimental trips (see below). All observed incidental interactions on a trip were assigned to the quarter when the vessel returned to port after completing the trip. Some quarterly estimates of interactions therefore involve interactions that occurred during an earlier quarter. Accordingly, these estimates are not the best source of information on seasonality of interactions.

The contractor's sampling records were used to approximate sampling probabilities. The sampling probabilities during the periods when additional (secondary) samples were drawn were computed by enumerating the number of call-ins during consecutive periods of comparable coverage. It was then assumed that the additional trips were selected with equal probability from those trips that had not been selected as part of the systematic sample. When coverage was below that of the anticipated systematic sample, the sampling probabilities were computed by enumerating all call-ins during this period and assuming that the trips sampled were selected with equal probability. Because the coverage level changed with the fluctuations in observer availability and fishing activity, trips were not selected with equal probability. Therefore, the Horvitz-Thompson estimator was used to estimate total interactions, as it takes into account unequal sampling probabilities. The incidental interaction records used to compute the Horvitz-Thompson estimator were those available in the Longline Observer Database System on 23 March 2007.

Confidence intervals for the quarterly incidental interactions were estimated using the approximated sampling probabilities and assuming that the number of incidental interactions per trip for a given species was an independent Poisson variate with a constant mean value. The assumption that the average rate of incidental interactions was constant throughout a quarter is questionable but necessary to compute confidence intervals. Confidence intervals for the yearly total were not computed, as it seems unreasonable to assume that incidental interaction rates were constant throughout the entire year.

During the third and fourth quarter of year 2005 and the first quarter of year 2006, several vessels participated in an experiment that involved alternating, within a set, between circle hooks and the hook style the vessel normally used. Additionally, in the third and fourth quarters of 2006, one vessel made seven trips as part of an experiment to test a deep setting fishing method. All trips involved in these experiments had an observer or scientist onboard. Since the protocol for these experiments fell under the current legal practices for this fishery, these trips were considered to be part of the Hawaii longline deep set longline fishery activity and the data were included in the estimation of interactions. Because the experimental trips had 100% coverage they were not part of the random sampling scheme. To estimate the total incidental interactions for all deep set longline fishing activity, the numbers of interactions observed on the experimental trips were added to the estimated numbers of interactions on trips subject to the random sampling scheme; i.e., all trips not participating in the experiments.

Additional estimates of marine mammal interactions required for stock assessment and management will be provided later in a separate report.

Table 1. Point estimates of the number of incidental interactions by species and corresponding 95% confidence intervals (C.I.) for the Hawaii deep set longline fishery in 2006. All protected species with an observed interaction are listed as well as species that most commonly interact with the fishery.

Species	Quarter								Annual Total
	1		2		3		4		
	Number of Incidental Interactions								
	Point Estimate	C.I.	Point Estimate	C.I.	Point Estimate	C.I.	Point Estimate	C.I.	Point Estimate
Turtles									
Loggerhead	0	[0,43]	0	[0,11]	0	[0,15]	0	[0,16]	0
Leatherback	0	[0,43]	2	[1,13]	7	[1,24]	0	[0,16]	9
Olive Ridley	6	[1,51]	18	[4,36]	30	[6,58]	0	[0,16]	54
Green	0	[0,43]	2	[1,13]	4	[1,21]	0	[0,16]	6
Albatrosses									
Black-footed	28	[5,112]	21	[8,39]	8	[2,26]	13	[2,34]	70
Laysan	0	[0,43]	0	[0,11]	7	[1,24]	0	[0,16]	7
Dolphins									
Striped	0	[0,43]	0	[0,11]	0	[0,15]	6	[1,25]	6
Bottlenose	1	[1,44]	0	[0,11]	0	[0,15]	0	[0,16]	1
Risso	0	[0,43]	5	[2,18]	0	[0,15]	0	[0,16]	5
Unidentified Dolphin	7	[1,64]	2	[1,13]	0	[0,15]	0	[0,16]	9
Whales									
Pilot	1	[1,44]	5	[1,18]	0	[0,15]	0	[0,16]	6
Humpback	0	[0,43]	0	[0,11]	0	[0,15]	0	[0,16]	0
False Killer	6	[1,51]	2	[1,13]	3	[1,19]	6	[1,25]	17
Beaked	0	[0,43]	0	[0,11]	0	[0,15]	0	[0,16]	0
Unidentified whale	11	[1,90]	0	[0,11]	0	[0,15]	3	[1,20]	14