

AGGREGATION AND OTHER REDUCTIONS FOR SEMI-SEPARABLE

MARKOV DECISION PROCESSES: THE LINEAR CASE

Roy Mendelsohn
Southwest Fisheries Center
National Marine Fisheries Service, NOAA
Honolulu, HI 96812

February 1978

DRAFT FOR COMMENT

I. Introduction

In a series of recent papers, I show how a semi-separable Markov decision process can be reduced to a problem with n constraints (the number of states) and n lower bounds. If $x \in X$ is a state, and $y \in X$ is a feasible decision, then a process is "semi-separable" (Mendelsohn 1978a)

$$P_{x,j}^y = P_j^y \quad \text{for all } x, j \in X; \text{ and all } y$$

where $P_{x,j}^y$ is the transition probability to state j . Let f_0, f_1, \dots, f_n be the optimal values of the Markov process, and let $v_0 = f_0$; $v_1 = f_1 - f_0, \dots, v_n = f_n - f_{n-1}$. Then I show the LP that solves the problem can be formulated as:

$$\begin{aligned} \min \quad & \sum_{x=0}^n (n+1-x)v_x \\ \text{s.t.} \quad & Av \geq b \\ & v_i \geq b_j^* \end{aligned} \tag{1.1}$$

where the $j+1$ element of b is the return from remaining in state j , b_j^* are simply determined lower bounds, and the j^{th} row of A is:

$$\left(1-\alpha, 1-\alpha \sum_{i=1}^n P_i^j, \dots, 1-\alpha \sum_{i=j}^n P_i^j, -\alpha \sum_{i=j+1}^n P_i^j, \dots, -\alpha P_n^j \right).$$

(see Mendelsohn 1978a for details). In Mendelsohn (1978b), I use the structure of this reduced LP to show how to easily recover an optimal policy from a solution to (1.1), and also to describe qualitative properties of an optimal policy for a wide class of semi-separable processes.

In this paper, I focus on a special case; where the decision set $Y(x)$ for each $x \in X$ is defined as:

$$Y(x) \equiv \{y : 0 \leq y \leq x\} \quad (1.2)$$

and the return is linear, that is:

$$G(x, y) = h \cdot (x-y); h \geq 0 \quad (1.3)$$

This is an example of a separable problem considered by Denardo (1968). However, (1.1) is a smaller problem than the equivalent problem in Denardo; and can be used to obtain even greater reductions in problem size.

In Mendelsohn (1978b), I show that a separable problem satisfying (1.2) and (1.3) always has a base stock policy optimal, if an optimal policy exists. Also, in the linear case, each b_j^* is identically h . the lower bounds can be removed by substituting $v_i' = v_i - h$ for $i = 1, \dots, n$. Before stating theorem 1, a lemma is needed.

Lemma 1.1 Let x be a nonnegative random variable defined on $[0, b]$, where $b \leq \infty$, and $F(x)$ its cdf. Then:

$$E(x) = \int_0^b [1 - F(x)] dx$$

Proof. See Karlin and Taylor (1975), p. 38.

□

Theorem 1.1 Assumption (1.2) and (1.3) imply the LP given (1.1) is equivalent to:

$$\begin{aligned} & \text{minimize } \sum_{x=0}^n v'_x \\ & \text{s.t. } Av \geq b \end{aligned} \tag{1.4}$$

where $b^i = \alpha E(s[i]) - (i+1)h$; and $s[i]$ is the transition function from state i .

Proof. By the substitution, $v_i = v'_i + h$. At i , this gives $A^i v$ and terms multiplies by h .

$$h \cdot \left((i+1) - \alpha \left(1 + \left(1 - p_0^i \right) + \left(1 - p_0^i - p_1^i \right) + \dots + \left(1 - p_0^i - p_1^i - \dots - p_n^i \right) \right) \right)$$

The inner terms are just $(i+1) - \alpha \sum_{i=0}^n (1 - F(i)) = (i+1) - \alpha E(s[i])$
by lemma 1.1.

□

It is interesting to note that the $(i+1)^{\text{st}}$ RHS is greater than the i^{th} , if and only if:

$$E(s[i+1]) - E(s[i]) \geq \frac{h}{\alpha} \tag{1.5}$$

In a continuous state version, this is equivalent to:

$$\frac{d}{di} (Es(i)) \geq \frac{1}{\alpha} \tag{1.6}$$

Equality in (1.6) is the two-period base stock size. In each row i of A , v^i is added to the row before. The algorithm checks that this "weighted derivative" is greater or less than the change in the two-period result. A related result can be found in Mendelssohn and Sobel (1977).

II. Reducing the Size of the LP

For many problems, even (1.4) will be a very large LP. Luckily, it is possible to greatly reduce (1.4). It is convenient for this discussion to let $b \equiv 0$, and retain the lower bounds $v_i \geq h$.

Often it is known a priori that the base stock size lies in the interval (j^L, j^u) . This suggests it may be possible to aggregate all the variables below j^L and all the variables above j^u . Theorem 2.1 states that the variables above j^u may be aggregated without loss of optimality; the variables below j^L may be aggregated but possibly with loss of optimality. However, the base stock size will still be in (j^L, j^u) .

Aggregate variables by letting:

$$c^u = \frac{1}{n-j^u+1} \left(\sum_{i=j^u}^n (n+1-i) \right); \quad c^L = \sum_{i=0}^{j^L} w^i (n+1-i)$$

where $w^k = \left(\left(1 - \alpha \sum_{i=k}^n p_i^{j^L} \right) / \sum_{i=0}^{j^L} \left(1 - \alpha \sum_{k=i}^n p_k^{j^L} \right) \right)$

and let $q_i^u = \frac{1}{n+1} \left(\sum_{k=j^u}^n a_k^i \right); \quad q_i^L = \sum_{k=0}^{j^L} \left(w^k a_k^i \right); \quad i = j^L, j^L+1, \dots, j^u$

where a_k^i is the (i, k) element of A . Let $\bar{v}^T = (v^L, v_{j^L+1}, \dots, v_{j^u-1}, v^u)$, and \bar{A} be A with rows 1 through j^L , j^u+1 through $n+1$ eliminated; columns 1 through j^L , j^u+1 through $n+1$ eliminated, and the first and last columns of \bar{A} are q^L and q^u , where $q^{L^T} = (q_{j^L}^L, q_{j^L+1}^L, \dots, q_{j^u}^L)$ and $q^{u^T} = (q_{j^L}^u, q_{j^L+1}^u, \dots, q_{j^u}^u)$.

Theorem 2.1 The reduced problem:

$$\min \sum_{i=j^L+1}^{j^u-1} (n+1-i)v_i + c^u v_u^u + c^L v^L \quad (1.7)$$

$$\text{s.t. } \bar{A}\bar{v} \geq 0$$

$$v_i \geq h, i = j^L+1, \dots, j^u-1; v^u \geq (n-j^u+1)h; v^L \geq (j^L+1)h$$

has the following properties:

- (i) The base stock size \bar{j} lies in (j^L, j^u) .
- (ii) There is no loss of optimality from aggregating $v_{j^u}, v_{j^u+1}, \dots, v_n$
- (iii) The loss of optimality from aggregating v_0, v_1, \dots, v_{j^L} is at most

$$\left[\max_{i=0,1,\dots,j^L} \left\{ (n+1-i) - \bar{u}A^i \right\} \right]^+ \frac{\alpha}{1-\alpha} hn,$$

where \bar{u} are the dual variables of the reduced problem.

Proof. A heuristic proof is offered for aggregating the upper variables, since a rigorous proof can be derived as a special case of the lower variables, which is more difficult. Since $j^* \in (j^L, j^u)$, each v_i , $i \geq j^u$,

must be identically h . Also, rows j^u+1 through $n+1$ cannot be tight, hence they can be eliminated. Averaging the transition is justified in that the new process, if it goes to a state greater than j^u , immediately goes at least to j^u (perhaps lower). These are the q_i^u 's.

For the lower variables, it is first necessary to show that j^L will not come out as the solution to (1.7). Suppose it does. The proof is to show that this implies $j^* \notin (j^L, j^u)$ in the original problem. Disaggregate by letting

$$v_i = w^i v^L \quad i = 0, \dots, j^L$$

$$v_i = \frac{1}{n-j^u+1} v^u \quad i = j^u, \dots, n+1$$

Then from proposition (1) in Zipkin the disaggregate solution is feasible in (1.1), and equality must hold in row j^L+1 , due to how the weights w^i were chosen.

Since v_{j^L+1}, \dots, v_n are at their lower bound and the coefficients on them are negative in that row, it is clear that an improved solution can only be obtained by decreasing v_0, v_1, \dots, v_{j^L} . This implies $v_{j^*-1}^*$ is at its lower bound, which contradicts the optimality of j^* .

Part (iii) is equation (4) in Zipkin (1977).

□

Theorem 2.2 suggests a method for finding an optimal policy. Aggregate together all variables with an index less than j^L , greater than j^u , and choose a partition on the remaining states, and aggregate up to each of those points. The following theorem shows that if the

base stock size found from this problem is at the variable representing the i^{th} set of aggregated states, then the true base stock size must lie in that partition.

Theorem 2.2 Assume $j^* \in (j^L, j^u)$. Divide (j^L, j^u) into intervals $(j^L, j^1]$, $(j^1, j^2]$, ..., (j^k, j^u) . Aggregate variables v^0, \dots, v_{j^L} and v_{j^u}, \dots, v_n as in theorem 2.1. Aggregate variables $v_{j^i}, v_{j^{i+1}}, \dots, v_{j^{i-1}}$ similarly to v_0, \dots, v_{j^L} , where the weights w^i are chosen from row j^i as in theorem 2.1. Then

- (i) An optimal base stock size for the aggregated problem lies in (j^L, j^u) .
- (ii) If j^i is an optimal base stock size for the aggregate problem, $j^* \in (j^i, j^{i+1})$.
- (iii) The error for the aggregate problem is at most

$$h \frac{\alpha n}{1-\alpha} \sum_{i=1}^K \left\{ \left[\max_{j=j^i, \dots, j^{i+1}-1} \left(n+1-(i+j) - \bar{u}A^{(i+j)} \right) \right]^+ \right\}$$

$$+ h \frac{\alpha n}{1-\alpha} \left(\left[\max_{i=0, \dots, j^L} (n+1-i - \bar{u}A^i) \right]^+ \right)$$

where \bar{u} is the vector of dual variables of the aggregated problem and A^i is the i^{th} column of A .

Proof. Part (iii) is again equation (4) in Zipkin (1977). If part (i) isn't true, then either the row equivalent to staying in j^L of j^u must be an equality. The contrapositive proof in this case is the same as in theorem 2.1.

Suppose j^i is the optimal base size for the aggregate problem. Assume $j^* < j^i$, then a similar contrapositive argument yields that j^* is not optimal in the original problem. If $j^* > j^{i+1}$, then set all the v_i 's, $i \geq j+1$ equal to h . Since this row is an inequality, it must be an equality in the original row. From the form of the disaggregated row, this can only come from changing $v_0, v_1, \dots, v_{j^{i+1}}$, so that $v_{j^*-1} = h$, which contradicts the optimality of j^* . □

An efficient search technique can now be devised. Aggregate to j^L and from j^u to $n+1$, and some small partition of (j^L, j^u) . If the partition is 10 more variables, this implies initially an LP that is $(12 \times 12)!$ Given the solution to this problem, reaggregate up to j^i and down to j^{i+1} , where j^i is the base stock size from the aggregate problem. Partition this interval as before, and repeat. This algorithm involves solving a series of extremely small LP's. The a posteriori error as given by (iii) can be used as a stopping rule for when it is no longer worth going further.

III. Finding the Interval (j^L, j^u)

Consider the problem:

$$f(x) = \max_{0 \leq y \leq x} \left\{ h \cdot (x-y) + \alpha E \left(f(s[y, D]) \right) \right\}$$

where D is a random variable, and it is assumed that $f(\cdot)$ and $s[\cdot, D]$ are differentiable (perhaps one-sided) and that $f(\cdot)$ is unimodal.

Assumptions that are sufficient for these conditions to be valid are given in Mendelssohn and Sobel (1977).

It is readily seen from that paper or the LP (1.1) that $f'(x) \geq h$. In fact, $f'(x) \equiv h$ if $x \geq y^*$, where y^* is an optimal base stock size. Therefore, if:

$$h \leq \alpha E\left\{h \cdot s^{[1]}[y, D]\right\} = \alpha h E\left\{s^{[1]}[y, D]\right\}$$

the base stock size must be no greater than y . Let y^* be the point where $E(s^{[1]}[y, D]) = \frac{1}{\alpha}$. Then the base stock size for the infinite horizon problem can be no greater than y^* . Let $j^u \equiv y^*$. If $s[y, D] = Ds[y]$, $\Pr\{D \geq 0\} = 1$, and $s[1]$ is continuous, it is easily seen that if \bar{y}^* is the solution to $s'[y] = \frac{1}{\alpha}$, then the infinite horizon problem base stock size is less than \bar{y}^* if $E(D) > 1$, is greater than \bar{y}^* if $E(D) < 1$, and is less than or equal to \bar{y}^* if $E(D) \equiv 1$.

Thus, in comparing the stochastic and deterministic problems, if \bar{y}^* is the deterministic solution, the stochastic solution is more conservative if and only if $E(s[\bar{y}^*, D]) - \bar{y}^*$ is a supermartingale. Using the deterministic solution as the base stock size is more conservative if $E(s[\bar{y}^*, D]) - \bar{y}^*$ is a submartingale. Unless $E(s[\bar{y}^*, D]) \gg \bar{y}^*$, it probably will not be worth it to calculate the stochastic optimum. Zipkin's (1977) results can be used to determine bounds on using this policy. However, particularly in situations where very low stock levels are highly undesirable, such as in managing renewable resources, using the deterministic solution as the base stock size to lower the risk of depletion will probably

outweigh the loss due to this bound. I conjecture that a similar result is valid when linear returns are combined with linear smoothing costs.

In a future paper, these results will be used to analyze several published models relating to managing stochastically varying fisheries.

Acknowledgment

I thank Professor Paul Zipkin of Columbia University for letting me use his unpublished results on aggregating linear programs.

References

- Karlin, S. and H. L. Taylor (1975) A first course in stochastic processes. (2d ed.) Academic Press, N.Y., 557 p.
- Mendelssohn, R. (1978a) Increasing computational efficiency for semi-separable Markov decision processes. U.S. Dep. Commer., Natl. Mar. Fish. Serv., SWFC Admin. Rep. 2H, 1978, 16 p.
- _____. (1978b) Recovering optimal policies and qualitative properties of optimal policies for semi-separable Markov decision processes. U.S. Dep. Commer., Natl. Mar. Fish. Serv., SWFC Admin. Rep. 3H, 1978, 7 p.
- Mendelssohn, R. and M. J. Sobel. (1977) Capital accumulations and the optimization of renewable resource models. Submitted to J. Econ. Theory.
- Zipkin, P. H. (1977) Bounds on the effect of aggregating variables in linear programs. Research paper #211, Graduate School of Business, Columbia University, N.Y., 29 p.